The effect of tobacco retailer density reduction on smoking in Belgium

Sarah Croes¹ • Leonor Gurariguata¹ • Brecht Devleesschauwer^{1,2} • Vanessa Gorasso¹

¹ Department of Epidemiology and Public Health, Sciensano, Brussels, Belgium • ² Department of Translational Physiology, Infectiology and Public Health, Ghent University, Ghent, Belgium





Smoking is one of the most significant risk factors for various diseases, posing a major **public health issue**.



Trends of current smoking suggest that tobacco control policies are needed to achieve targets.

The Belgian Tobacco-Free Generation **Plan** aims to reduce smoking by limiting access tobacco, including the reduction of tobacco selling points.

How much burden can be avoided, 🞾 evaluated by health impact assessment?



~ 1% disease burden avoided by reducing tobacco sell-points

We would like to express our gratitude to our colleagues at SPF Public Health and SEO for providing the data and sharing valuable insights. A special thank you to my wonderful colleagues from the Health Indicator Team, especially Sarah and Vanessa, for their assistance with the layout of this poster. This research was funded by the European Union through the Technical Support Instrument (TSI) project "Resources Hub for sustainable investing in Health" (2022-2024).



baseline scenario

Dark green = high density Light green = low density

shading

in baseline scenario